Brain Research, 612 (1993) 357–359
Elsevier Science Publishers B.V.

Author Index

Abell, C.W., see Westlund, K.N. (612) 221
Ahmed, I., see Aravich, P.F. (612) 180
Aravich, P.F., Rieg, T.S., Ahmed, I. and Lauterio, T.J.
Fluoxetine induces vasopressin and oxytocin abnormalities in food-restricted rats given voluntary exercise: relationship to anorexia nervosa (612) 180
Awiszus, F. and Schäfer, S.S.
Subdivision of primary afferents from passive cat muscle spindles based on a single slow-adaptation parameter (612) 110

Back, S.A., Colon, M., Fallon, J.H., Meyskens Jr., F.L. and Loughlin, S.E.
Ventral mesencephalic and cortical transplants into the rat striatum display enhanced activity for neutral endopeptidase 24.11 ('enkephalinase'; CALLA) (612) 85
Barbacid, M., see Steininger, T.L. (612) 330
Barber, D.J., see Cooper, S.J. (612) 313
Bartfai, T., see Girotti, P. (612) 258
Bedran de Castro, J.C., see Bedran de Castro, M.T.B. (612) 243
Bedran de Castro, M.T.B., Bedran de Castro, J.C. and Vanderlei Menani, J.
The effects of forebrain multiple lesions on the pressor response induced by bilateral carotid occlusion in conscious rats (612) 243
Behbehani, M.M., Liu, H., Jiang, M., Pun, R.Y.K. and Shipley, M.T.
Activation of serotonin$_{1A}$ receptors inhibits midbrain periaqueductal gray neurons of the rat (612) 56
Belizario, J., see Coceani, F. (612) 165
Bellgowan, P.S., see Helmstetter, F.J. (612) 253
Berger, P.J., Cooke, I.R.C., Soust, M. and Wilkinson, M.H.
Origin of tonic electrical activity patterns in the diaphragm of the fetal lamb (612) 299
Bertorelli, R., see Girotti, P. (612) 258
Bonavera, J.J., Kalra, S.P. and Kalra, P.S.
Mode of action of interleukin-1 in suppression of pituitary LH release in castrated male rats (612) 1
Brown, L.D., Kim, K.-M., Nakajima, Y. and Nakajima, S.
The role of G protein in muscarinic depolarization near resting potential in cultured hippocampal neurons (612) 200
Bryant, B.P., see Komai, M. (612) 122
Butcher, S.P., see Lawrence, J.A. (612) 326

Castillo, N., see Weiss, G. (612) 354
Chagoya de Sánchez, V., Hernández Muñoz, R., Suárez, J., Vidrio, S., Yáñez, L. and Muñoz, M.D.
Day–night variations of adenosine and its metabolizing enzymes in the brain cortex of the rat – possible physiological significance for the energetic homeostasis and the sleep–wake cycle (612) 115
Chambille, I., Doyle, S. and Servière, J.
Photic induction and circadian expression of Fos-like protein. Immunohistochemical study in the retina and suprachiasmatic nuclei of hamster (612) 138
Chugani, H.T., see Rocha, L. (612) 247
Clement, M.E. and McCall, R.B.
Lateral tegmental field involvement in the central sympathoinhibitory action of 8-OHDPAT (612) 78
Coceani, F., Lees, J., Mancilla, J., Belizario, J. and Dinarello, C.A.
Interleukin-6 and tumor necyosis factor in cerebrospinal fluid: changes during pyrogen fever (612) 165
Coenen, A.M.L., see Inoue, M. (612) 35
Colon, M., see Back, S.A. (612) 85
Consolo, S., see Girotti, P. (612) 258
Cooke, I.R.C., see Berger, P.J. (612) 299
Cooper, S.J. and Barber, D.J.
The benzodiazepine receptor partial agonist bretazenil and the partial inverse agonist Ro 15-4513: effects on salt preference and aversion in the rat (612) 313
Crockett, D.P., Maslany, S., Harris, S.L. and Egger, M.D.
Enhanced cytochrome-oxidase staining of the cuneate nucleus in the rat reveals a modifiable somatotopic map (612) 41

DeBold, J.F., see Frye, C.A. (612) 130
De Vries, M.J., Nunes Cardozo, B., Van der Want, J., De Wolf, A. and Meijer, J.H.
Glutamate immunoreactivity in terminals of the retinohypothalamic tract of the brown Norwegian rat (612) 231
De Wolf, A., see De Vries, M.J. (612) 231
Dinarello, C.A., see Coceani, F. (612) 165
Doyen, C., see Kidd, E.J. (612) 289
Doyle, S., see Chambille, I. (612) 138
Duysens, J., see Inoue, M. (612) 35

Egger, M.D., see Crockett, D.P. (612) 41
Engel Jr., J., see Rocha, L. (612) 247

Fabricius, M., Jensen, L.H. and Lauritzen, M.
Microdialysis of interstitial amino acids during spreading depression and anoxic depolarization in rat neocortex (612) 61
Fallon, J.H., see Back, S.A. (612) 85
Fattaccini, C.-M., see Kidd, E.J. (612) 289
Fernandez, M., see Weiss, G. (612) 354
Fisone, G., see Girotti, P. (612) 258
Fredholm, B.B., see Georgiev, V. (612) 271
Frye, C.A. and DeBold, J.F.
3α-OH-DHP and 5α-THDOC implants to the ventral tegmental area facilitate sexual receptivity in hamsters after progesterone priming to the ventral medial hypothalamus (612) 130
Fujii, M., see Koyama, S. (612) 345

Gai, W.-P., see Zhou, X.-F. (612) 322
Georgiev, V., Johansson, B. and Fredholm, B.B.
Long-term caffeine treatment leads to a decreased susceptibility to NMDA-induced clonic seizures in mice without changes in adenosine A_1 receptor number (612) 271
Girotti, P., Bertorelli, R., Fisone, G., Land, T., Langel, Ü., Consolo, S. and Bartfai, T.
N-terminal galanin fragments inhibit the hippocampal release of acetylcholine in vivo (612) 258
Görcs, T., see Töröcsik, A. (612) 306
Gozlan, H., see Kidd, E.J. (612) 289
Gurnett, C.A., Whiteis, C.A., Pardini, B.J. and Schmid Jr., P.G.
Norepinephrine release from guinea pig cardiac sympathetic nerves is insensitive to ryanodine under physiological conditions (612) 238

Hama, N., see Kaneko, T. (612) 104
Hamon, M., see Kidd, E.J. (612) 289
Harik, S.I., see Sullivan, H.C. (612) 336
Harris, S.L., see Crockett, D.P. (612) 41
Hazama, F., see Yamada, E. (612) 210
Heinemann, U., see Kral, T. (612) 278
Helmstetter, F.J. and Bellgowan, P.S.
Lesions of the amygdala block conditional hypoalgesia on the tail flick test (612) 253
Hernández Muñoz, R., see Chagoya de Sánchez, V. (612) 115
Hineno, T., see Mizobuchi, M. (612) 319
Hirai, O., see Kaneko, T. (612) 104
Hiratani, K., see Mizobuchi, M. (612) 319
Hirayama, T., see Koyama, S. (612) 345

Imura, H., see Kaneko, T. (612) 104
Inoue, M., Duysens, J., Vossen, J.M.H. and Coenen, A.M.L.
Thalamic multiple-unit activity underlying spike-wave discharges in anesthetized rats (612) 35

Jensen, L.H., see Fabricius, M. (612) 61
Jiang, M., see Behbehani, M.M. (612) 56
Joëls, M., see Karst, H. (612) 172
Johansson, B., see Georgiev, V. (612) 271
Jouvet, M., see Sei, H. (612) 351

Kakimoto, Y., see Mizobuchi, M. (612) 319
Kalra, P.S., see Bonavera, J.J. (612) 1
Kalra, S.P., see Bonavera, J.J. (612) 1
Kaneko, T., Shirakami, G., Nakao, K., Nagata, I., Nakagawa, O., Hama, N., Suga,

S.-i., Miyamoto, S., Kubo, H., Hirai, O., Kikuchi, H. and Imura, H.
C-type natriuretic peptide (CNP) is the major natriuretic peptide in human cerebrospinal fluid (612) 104
Karnaze, D., see Weiss, G. (612) 354
Karst, H., Wadman, W.J. and Joëls, M.
Long-term control by corticosteroids of the inward rectifier in rat CA1 pyramidal neurons, in vitro (612) 172
Katayama, Y., see Koyama, S. (612) 345
Kelly, J.S., see Lawrence, J.A. (612) 326
Kemenes, G., Zakharov, I.S., Vehovszky, Á. and S.-Rózsa, K.
Neurophysiological correlates of tactile stimulus-induced whole-body eversion, a novel type of behavior in the snail *Helix pomatia* L. (612) 16
Kidd, E.J., Laporte, A.M., Langlois, X., Fattaccini, C.-M., Doyen, C., Lombard, M.C., Gozlan, H. and Hamon, M.
5-HT_3 receptors in the rat central nervous system are mainly located on nerve fibres and terminals (612) 289
Kikuchi, H., see Kaneko, T. (612) 104
Kim, K.-M., see Brown, L.D. (612) 200
Kim, M. and Mistretta, C.M.
4-Aminopyridine reduces chorda tympani nerve taste responses to potassium and alkali salts in rat (612) 96
Kishi, K., see Kuroda, M. (612) 339
Klein, R., see Steininger, T.L. (612) 330
Kogure, K., see Tomioka, C. (612) 216
Köller, H., Siebler, M., Pekel, M. and Müller, H.W.
Depolarization of cultured astrocytes by leukotriene B_4. Evidence for the induction of a K^+ conductance inhibitor (612) 28
Komai, M. and Bryant, B.P.
Acetazolamide specifically inhibits lingual trigeminal nerve responses to carbon dioxide (612) 122
Koyama, S., Katayama, Y., Maejima, S., Hirayama, T., Fujii, M. and Tsubokawa, T.
Thalamic neuronal hyperactivity following transection of the spinothalamic tract in the cat: involvement of *N*-methyl-D-aspartate receptor (612) 345
Krakower, T.J., see Westlund, K.N. (612) 221
Kral, T., Luhmann, H.J., Mittmann, T. and Heinemann, U.
Role of NMDA receptors and voltage-activated calcium channels in an in vitro model of cerebral ischemia (612) 278
Kubo, H., see Kaneko, T. (612) 104
Kuroda, M., Murakami, K., Oda, S., Shinkai, M. and Kishi, K.
Direct synaptic connections between thalamocortical axon terminals from the mediodorsal thalamic nucleus (MD) and corticothalamic neurons to MD in the prefrontal cortex (612) 339
Kwan, S.-W., see Westlund, K.N. (612) 221

Land, T., see Girotti, P. (612) 258
Langel, Ü., see Girotti, P. (612) 258
Langlois, X., see Kidd, E.J. (612) 289
Laporte, A.M., see Kidd, E.J. (612) 289
Lauritzen, M., see Fabricius, M. (612) 61
Lauterio, T.J., see Aravich, P.F. (612) 180
Lawrence, J.A., Olverman, H.J., Shirakawa, K., Kelly, J.S. and Butcher, S.P.
Binding of 5-HT_{1A} receptor and 5-HT transporter ligands in rat cortex and hippocampus following cholinergic and serotonergic lesions (612) 326
Leach, M.J., see Lees, G. (612) 190
Lees, G. and Leach, M.J.
Studies on the mechanism of action of the novel anticonvulsant lamotrigine (Lamictal) using primary neuroglial cultures from rat cortex (612) 190
Lees, J., see Coceani, F. (612) 165
Liu, H., see Behbehani, M.M. (612) 56
Lombard, M.C., see Kidd, E.J. (612) 289
Loughlin, S.E., see Back, S.A. (612) 85
Lucero, K., see Weiss, G. (612) 354
Luhmann, H.J., see Kral, T. (612) 278

Maejima, S., see Koyama, S. (612) 345
Mancilla, J., see Coceani, F. (612) 165
Maslany, S., see Crockett, D.P. (612) 41
McCall, R.B., see Clement, M.E. (612) 78
Meijer, J.H., see De Vries, M.J. (612) 231
Meyskens Jr., F.L., see Back, S.A. (612) 85
Mistretta, C.M., see Kim, M. (612) 96
Mittmann, T., see Kral, T. (612) 278
Miyamoto, S., see Kaneko, T. (612) 104
Mizobuchi, M., Hineno, T., Kakimoto, Y. and Hiratani, K.
Increase of plasma adrenocorticotrophin and cortisol in 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)-treated dogs (612) 319
Müller, H.W., see Köller, H. (612) 28
Muñoz, M.D., see Chagoya de Sánchez, V. (612) 115
Murakami, K., see Kuroda, M. (612) 339

Nagata, I., see Kaneko, T. (612) 104
Nakagawa, O., see Kaneko, T. (612) 104
Nakajima, S., see Brown, L.D. (612) 200
Nakajima, Y., see Brown, L.D. (612) 200
Nakao, K., see Kaneko, T. (612) 104
Nishioka, K., see Tomioka, C. (612) 216
North, R.A., see Sugita, S. (612) 151
Nunes Cardozo, B., see De Vries, M.J. (612) 231

Oda, S., see Kuroda, M. (612) 339
Olverman, H.J., see Lawrence, J.A. (612) 326

Palfrey, H.C., see Steininger, T.L. (612) 330
Pardini, B.J., see Gurnett, C.A. (612) 238
Pekel, M., see Köller, H. (612) 28
Penev, P.D., Turek, F.W. and Zee, P.C.
Monoamine depletion alters the entrainment and the response to light of the circadian activity rhythm in hamsters (612) 156
Pun, R.Y.K., see Behbehani, M.M. (612) 56

Rakovska, A., see Töröcsik, A. (612) 306
Randich, A., see Thurston, C.L. (612) 70
Rieg, T.S., see Aravich, P.F. (612) 180
Rocha, L., Tatsukawa, K., Chugani, H.T. and Engel Jr., J.
Benzodiazepine receptor binding following chronic treatment with naloxone, morphine and met-enkephalin in normal rats (612) 247
Rush, R.A., see Zhou, X.-F. (612) 322

S.-Rózsa, K., see Kemenes, G. (612) 16
Saito, T.
4-Aminopyridine causes reduction of cytochromes partly originated in glia with enhanced evoked potentials in the olfactory cortex slice (612) 9
Sakai, K., see Sei, H. (612) 351
Salamone, J.D., see Steinpreis, R.E. (612) 263
Schäfer, S.S., see Awiszus, F. (612) 110
Schmid Jr., P.G., see Gurnett, C.A. (612) 238
Sei, H., Sakai, K., Yamamoto, M. and Jouvet, M.
Spectral analyses of PGO-on neurons during paradoxical sleep in freely moving cats (612) 351
Servière, J., see Chambille, I. (612) 138
Shinkai, M., see Kuroda, M. (612) 339
Shipley, M.T., see Behbehani, M.M. (612) 56
Shirakami, G., see Kaneko, T. (612) 104
Shirakawa, K., see Lawrence, J.A. (612) 326
Siebler, M., see Köller, H. (612) 28
Soust, M., see Berger, P.J. (612) 299
Starnes, A., see Thurston, C.L. (612) 70
Steininger, T.L., Wainer, B.H., Klein, R., Barbacid, M. and Palfrey, H.C.
High-affinity nerve growth factor receptor (Trk) immunoreactivity is localized in cholinergic neurons of the basal forebrain and striatum in the adult rat brain (612) 330
Steinpreis, R.E. and Salamone, J.D.
The role of nucleus accumbens dopamine in the neurochemical and behavioral effects of phencyclidine: a microdialysis and behavioral study (612) 263
Suárez, J., see Chagoya de Sánchez, V. (612) 115
Suga, S.-i., see Kaneko, T. (612) 104
Sugita, S. and North, R.A.
Opioid actions on neurons of rat lateral amygdala in vitro (612) 151
Sullivan, H.C. and Harik, S.I.
ATP-sensitive potassium channels are not expressed in brain microvessels (612) 336

Tatsukawa, K., see Rocha, L. (612) 247
Thurston, C.L., Starnes, A. and Randich, A.
Changes in nociception, arterial blood pressure and heart rate produced by intravenous morphine in the conscious rat (612) 70
Tomioka, C., Nishioka, K. and Kogure, K.
A comparison of induced heat-shock protein in neurons destined to survive and those destined to die after transient ischemia in rats (612) 216
Töröcsik, A., Rakovska, A., Görcs, T. and Vizi, E.S.

Effect of neurotensin and immunneutralization with anti-neurotensin-serum on dopaminergic–cholinergic interaction in the striatum (612) 306
Tsubokawa, T., see Koyama, S. (612) 345
Turek, F.W., see Penev, P.D. (612) 156

Vanderlei Menani, J., see Bedran de Castro, M.T.B. (612) 243
Van der Want, J., see De Vries, M.J. (612) 231
Vehovszky, Á., see Kemenes, G. (612) 16
Vidrio, S., see Chagoya de Sánchez, V. (612) 115
Vizi, E.S., see Töröcsik, A. (612) 306
Vossen, J.M.H., see Inoue, M. (612) 35

Wadman, W.J., see Karst, H. (612) 172
Wainer, B.H., see Steininger, T.L. (612) 330
Weiss, G., Lucero, K., Fernandez, M., Karnaze, D. and Castillo, N.
The effect of adrenalectomy on the circadian variation in the rate of kindled seizure development (612) 354
Westlund, K.N., Krakower, T.J., Kwan, S.-W. and Abell, C.W.
Intracellular distribution of monoamine oxidase A in selected regions of rat and monkey brain and spinal cord (612) 221
Whiteis, C.A., see Gurnett, C.A. (612) 238
Wilkinson, M.H., see Berger, P.J. (612) 299

Yamada, E. and Hazama, F.
Different stability of neurofilaments for trypsin treatment after axotomy in the dorsal motor nucleus of the vagal nerve and the hypoglossal nucleus (612) 210
Yamamoto, M., see Sei, H. (612) 351
Yáñez, L., see Chagoya de Sánchez, V. (612) 115

Zakharov, I.S., see Kemenes, G. (612) 16
Zee, P.C., see Penev, P.D. (612) 156
Zhou, X.-F., Gai, W.-P. and Rush, R.A.
CGRP immunoreactive neurons in rat dorsal root ganglia do not all contain low-affinity NGF receptor immunoreactivity (612) 322